

National Immunisation Program Schedule

From 1 July 2018

Age	Disease	Vaccine brand
Adolescent vaccination (also see influenza vaccine)		
10-<15 years (School programs ^d)	<ul style="list-style-type: none"> Human papillomavirus (HPV)^e Diphtheria, tetanus, pertussis (whooping cough) 	Gardasil [®] 9 Boostrix [®]
Adult vaccination (also see influenza vaccine)		
15–49 years Aboriginal and Torres Strait Islander people with medical risk factors ^c	<ul style="list-style-type: none"> Pneumococcal 	Pneumovax 23 [®]
50 years and over Aboriginal and Torres Strait Islander people	<ul style="list-style-type: none"> Pneumococcal 	Pneumovax 23 [®]
65 years and over	<ul style="list-style-type: none"> Pneumococcal 	Pneumovax 23 [®]
70–79 years^f	<ul style="list-style-type: none"> Shingles (herpes zoster) 	Zostavax [®]
Pregnant women	<ul style="list-style-type: none"> Pertussis (whooping cough)^g Influenza^h 	Boostrix [®] or Adacel [®]

Funded annual influenza vaccination^h

6 months and over with certain medical risk factors^a

Aboriginal and Torres Strait Islander children 6 months to less than 5 years

Aboriginal and Torres Strait Islander people 15 years and over

65 years and over

Pregnant women

^a Hepatitis B vaccine: Should be given to all infants as soon as practicable after birth. The greatest benefit is if given within 24 hours, and must be given within 7 days.

^b Rotavirus vaccine: First dose must be given by 14 weeks of age, the second dose by 24 weeks of age.

^c Refer to the current edition of *The Australian Immunisation Handbook* for all medical risk factors.

^d Contact your state or territory health service for school grades eligible for vaccination.

^e Observe Gardasil[®]9 dosing schedules by age and at-risk conditions. 2 doses: 9 to <15 years - 6 months minimum interval. 3 doses: ≥15 years and/or have certain medical conditions - 0, 2 and 6 month schedule. Only 2 doses funded on the NIP unless 12-13 year old has certain medical risk factors.

^f All people aged 70 years old, with a five year catch-up program for people aged 71-79 years old until 31 October 2021.

^g Single dose recommended each pregnancy, ideally between 28-32 weeks, but may be given up until delivery.

^h Refer to annual influenza information for recommended vaccine brand for age.

- Contact your State and Territory Health Department for further information on any additional immunisation programs specific to your State or Territory.

- All people aged less than 20 years are eligible for free catch up vaccines.

- Adult refugees and humanitarian entrants are eligible for free catch up vaccines.

For more information

health.gov.au/immunisation

State/Territory

Australian Capital Territory
New South Wales
Northern Territory
Queensland
South Australia
Tasmania
Victoria
Western Australia

Contact Number

(02) 6205 2300
1300 066 055
(08) 8922 8044
13 HEALTH (13 4325 84)
1300 232 272
1800 671 738
1300 882 008
(08) 9321 1312